

Meliosma lanceolata 樹皮から単離した
トリテルペン配糖体

阿部フミ子*、山内辰郎*、澁谷博孝、北川 勲**

Phytochemistry, 42(3), 809-814(1996)

Triterpenoid Glycosides from Bark of
Meliosma lanceolata

Fumiko Abe*, Tatsuo Yamauchi*, Hirotaka Shibuya,
and Isao Kitagawa**

ABSTRACT From the bark of *Meliosma lanceolata*, nine triterpenoid glycosides including the 28-O- β -D-glucopyranosyl ester of bayogenin-3-O- β -D-glucuronopyranoside, its 4'-O- β -D-galactopyranoside and 4'-O- α -L-arabinopyranoside were isolated and the structures characterized along with that of 28-O- β -D-glucopyranosyl-bayogenin-3-O- β -D-4'-anhydro-4',5'-didehydroglucuronopyranoside. Bisdesmosidic triosides of hederagenin were obtained as minor components.

抄録 インドネシア・スマトラ島で採集した*Meliosma lanceolata* 樹皮から9種のトリテルペン配糖体を単離し、それらの化学構造を解明した。

* Faculty of Pharmaceutical Sciences, Fukuoka University

福岡大学薬学部

** Faculty of Pharmaceutical Sciences, Kinki University

近畿大学薬学部